



TABLE TO PREPARE POLYACRYLAMIDE GELS FOR NONDENATURING DNA ELECTROPHORESIS

Acrylamide concentration in the gel		3.5%	5.0%	8.0%	12.0%	15.0%	20.0%
Description	Catalog No	Reagents to prepare 100.0 ml working solution					
30% Acrylamide/Bisacrylamide Solution (19:1)	A00008	11.7 ml	16.7 ml	26.7 ml	40.0 ml	50.0 ml	66.7 ml
TBE (10x) Buffer	T00011	10.0 ml	10.0 ml	10.0 ml	10.0 ml	10.0 ml	10.0 ml
Distilled H ₂ O		77.6 ml	72.6 ml	62.6 ml	49.3 ml	39.3 ml	22.6 ml
<i>Mix thoroughly ^(*), then add</i>							
10% Ammonium Persulfate	} G00001	0.7 ml	0.7 ml	0.7 ml	0.7 ml	0.7 ml	0.7 ml
TEMED		35 µl	35 µl	35 µl	35 µl	35 µl	35 µl
<i>Mix the working solution by gentle swirling. Cast the gel, allow to polymerize for one hour.</i>							

(*) Deaeration of the solution at this point optimizes the polymerization of the gel and increases the repeatability of the results.

!! ENVIRONMENTAL PROTECTION NOTICE !!

In order to minimize the release of acrylamide to the environment, please polymerize any unused amount of monomer solution (acrylamide/bisacrylamide) according to the above table and the common practice. Let the solution to polymerize for one hour or more. Discard the formed gel.